

Amazon Tipping Point

In the 1970s, Brazilian scientist Eneas Salati shattered the long held dogma that vegetation is simply the consequence of climate and has no influence on climate whatsoever (1). Using isotopic ratios of oxygen in rainwater samples collected from the Atlantic to the Peruvian border, he was able to demonstrate unequivocally that the Amazon generates approximately half of its own rainfall by recycling moisture 5 to 6 times as airmasses move from the Atlantic across the basin to the west.

From the start, the demonstration of the hydrological cycle of the Amazon raised the question of how much deforestation would be required to cause the cycle to degrade to the point of being unable to support rain forest ecosystems.

High levels of evaporation and transpiration that forests produce throughout the year contribute to a wetter atmospheric boundary layer than would be the case with non-forest. This surface-atmosphere coupling is more important where large-scale factors for rainfall formation are weaker, such as in central and eastern Amazonia. Near the Andes, the impact of at least modest deforestation is less dramatic because the general ascending motion of airmasses in this area induces high levels of rainfall in addition to that expected from local evaporation and transpiration.

Where might the tipping point be for deforestation-generated degradation of the hydrological cycle? The very first model to examine this question (2) showed that at about 40% deforestation, central, southern and eastern Amazonia would experience diminished rainfall and a lengthier dry season, predicting a shift to savanna vegetation to the east.

Moisture from the Amazon is important to rainfall and human wellbeing because it contributes to winter rainfall for parts of the La Plata basin, especially southern Paraguay, southern Brazil, Uruguay and central-eastern Argentina; in other regions, the moisture passes over the area, but does not precipitate out. Although the amount contributing to rainfall in southeastern Brazil is smaller than in other areas, even small amounts can be a welcome addition to urban reservoirs.

The importance of Amazon moisture for Brazilian agriculture south of the Amazon is complex but not trivial. Perhaps most important is the partial contribution of dry season Amazon evapotranspiration to rainfall in southeastern South America. Forests maintain an evapotranspiration rate year-round, whereas evapotranspiration in pastures

is dramatically lower in the dry season. As a consequence, models suggest a longer dry season after deforestation.

In recent decades, new forcing factors have impinged on the hydrological cycle: climate change and widespread use of fire to eliminate felled trees and clear weedy vegetation. Many studies show that in the absence of other contributing factors, 4 degrees Celsius of global warming would be the tipping point to degraded savannas in most of the central, southern, and eastern Amazon. Widespread use of fire leads to drying of surrounding forest and greater vulnerability to fire in the subsequent year.

We believe that negative synergies between deforestation, climate change, and widespread use of fire indicate a tipping point for the Amazon system to flip to non-forest ecosystems in eastern, southern and central Amazonia at 20–25% deforestation.

The severity of the droughts of 2005, 2010 and 2015–16 could well represent the first flickers of this ecological tipping point. These events, together with the severe floods of 2009, 2012 (and 2014 over SW Amazonia), suggest that the whole system is oscillating. For the last two decades the dry season over the southern and eastern Amazon has been increasing. Large scale factors such as warmer sea surface temperatures over the tropical North Atlantic also seem to be associated with the changes on land.

We believe that the sensible course is not only to strictly curb further deforestation, but also to build back a margin of safety against the Amazon tipping point, by reducing the deforested area to less than 20%, for the commonsense reason that there is no point in discovering the precise tipping point by tipping it. At the 2015 Paris Conference of the Parties, Brazil committed to 12 million ha of reforestation by 2030. Much or most of this reforestation should be in southern and eastern Amazonia. The hydrological cycle of the Amazon is fundamental to human wellbeing in Brazil and adjacent South America.

– Thomas E. Lovejoy and Carlos Nobre

REFERENCES

1. E. Salati, A. Dall'Olio, E. Matsui, J. R. Gat, Recycling of Water in the Amazon, Brazil: an isotopic study. *Water Resour. Res.* **15**, 1250–1258 (1979).
2. G. Sampaio, C. A. Nobre, M. H. Costa, P. Satyamurty, B. S. Soares-Filho, M. Cardoso, Regional climate change over eastern Amazonia caused by pasture and soybean cropland expansion. *Geophys. Res. Lett.* **34**, L17709 (2007).

10.1126/sciadv.aat2340

Citation: T. E. Lovejoy, C. Nobre, Amazon Tipping Point. *Sci. Adv.* **4**, eaat2340 (2018).



Thomas E. Lovejoy is University Professor in the Department of Environmental Science and Policy at George Mason University. Email: tlovejoy@unfoundation.org



Carlos Nobre is a Member of the Brazilian Academy of Sciences and Senior Fellow of World Resources Institute Brazil.

Copyright © 2018
The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works. Distributed under a Creative Commons Attribution NonCommercial License 4.0 (CC BY-NC).

Amazon Tipping Point

Thomas E. Lovejoy and Carlos Nobre

Sci Adv 4 (2), eaat2340.
DOI: 10.1126/sciadv.aat2340

ARTICLE TOOLS <http://advances.sciencemag.org/content/4/2/eaat2340>

REFERENCES This article cites 2 articles, 0 of which you can access for free
<http://advances.sciencemag.org/content/4/2/eaat2340#BIBL>

PERMISSIONS <http://www.sciencemag.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of Service](#)

Science Advances (ISSN 2375-2548) is published by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. 2017 © The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works. The title *Science Advances* is a registered trademark of AAAS.